Advances in Agricultural Biotechnology

ISSN: 3117-6631
https://inkwise.org/index.php/AAB

Inkwise Press

Pilot Study Open Access

# Insect Protein Production From Agri-Food Side Streams: Pilot Study of Substrate Variability

**Ilze-Mara Bebris[1*], Ceda Erakovic[2] and Denijs Gheysen-Del[3]**

[1]Centre for Insect Biotechnology, Talsi Centre for Plant Genome Engineering, Talsi, Latvia
[2]Department of Bioprocess Engineering, Banat Institute of Agricultural Biotechnology, Zrenjanin, Serbia
[3]Department of Agricultural Economics, Pays des Collines Centre for Plant Genome Engineering, Frasnes, Belgium

*Correspondence to: Ilze-Mara Bebris, Centre for Insect Biotechnology, Talsi Centre for Plant Genome Engineering, Talsi, Latvia, E-mail: i.bebris@tcpge.lv

Received: 01 May 2023; Accepted: 09 August 2023; Published: 06 January 2024

Copyright: © 2024 Ilze-Mara Bebris, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

*There is real interest in insect protein production from agri-food side streams right now, and real uncertainty about exactly where it delivers the most value. We pull together the current evidence for insect protein production from agri-food side streams, covering the methodological, practical, regulatory and cost angles, with an eye on where practice has genuinely settled and where researchers and regulators are still largely working it out for themselves. Selective, context-specific adoption looks better supported by the evidence gathered here than either broad rollout or blanket avoidance.*

**Keywords:** Insect Protein Production From Agri-Food Side Streams; Agricultural biotechnology; Gene editing; Evidence synthesis; Biosafety regulation

## Introduction

Insect Protein Production From Agri-Food Side Streams has moved from a proof-of-concept curiosity to something a breeding or production programme can actually put to use, and that shift has outpaced the evidence base that is supposed to guide how it gets deployed.[1] The present pilot study sets out what the evidence currently says about insect protein production from agri-food side streams, where the science and the regulatory pathway have genuinely converged, and where practice is still mostly improvised.[2]

Interest in insect protein production from agri-food side streams has grown quickly as the underlying molecular tools have become reliable and affordable enough to move out of a specialist lab and into an ordinary breeding or production pipeline -- but that speed has left real variation in how different teams define, measure and report on it.[3]

We write this from the standpoint of someone deciding whether insect protein production from agri-food side streams is worth pursuing for their own programme, not as a purely academic exercise, and we come back to that practical question throughout.

## Related Work

The published work on insect protein production from agri-food side streams makes a reasonably convincing case for benefit overall, though that case is considerably stronger for some of its sub-questions than others.[4]

Early work in this area was mostly proof-of-concept -- a handful of edited lines, one growth cycle, a promising result. What has followed since has been slower and more methodical: replication, off-target screening, and a small but growing number of studies that actually track outcomes across multiple generations or seasons rather than one.[5]

## Methodology and Field Approach

Some parts of the insect protein production from agri-food side streams evidence base are backed by several agreeing independent studies; other parts rest on one or two single-lab reports whose findings may or may not travel to a different genetic background, species or growing environment.[6][7] Where the evidence is solid, practice has started to converge; where it is not, breeders and regulators are still working it out case by case, understandably.[8]

A theme that keeps coming up in the insect protein production from agri-food side streams literature is the gap between what works under controlled lab or greenhouse conditions and what has actually been shown to hold up once a line moves into confined or open field trials, with their usual constraints on time, containment and cost.[9]

Comparing results across the available insect protein production from agri-food side streams studies is genuinely difficult -- different genetic backgrounds, different editing constructs, different ways of defining a successful outcome -- so the pooled figures reported in this review are best read as a rough indication rather than a precise number.[10][11]

| Question | Current evidence | Practical implication |
|---|---|---|
| Strength of supporting evidence | Uneven across sub-questions | Weight recommendations by evidence tier |
| Consistency across genetic backgrounds | Variable transfer reported | Local validation before wider rollout |
| Metrics used | Mixed molecular and field metrics | Prefer studies with real field-performance data |
| Cost / resource implications | Rarely reported in detail | Assess local cost-effectiveness separately |

Table 1: Summary of key considerations for insect protein production from agri-food side streams.

## Evaluation Considerations Across Studies

Study quality across the insect protein production from agri-food side streams literature is mixed. A minority of the work is properly powered with a real unedited-control comparison; a larger share reports results from a single line, a single trial, with no formal control -- useful as a signal, but not something a breeding programme would want to commit resources to on that basis alone.[12]

How "success" gets measured for insect protein production from agri-food side streams also varies a lot between studies, and a result measured against a molecular or lab benchmark is simply not the same thing as one measured against field performance or downstream product quality, even when both get described in similar language in the abstract.[13]

## Implementation Considerations

Getting insect protein production from agri-food side streams to actually work at production scale involves more than the core molecular method: reliable regeneration of edited material into viable plants or animals, a way to confirm the edit and screen for off-target effects, and a plan for what to do when a line underperforms all matter as much as the underlying technique.[14] More than one of the programmes we reviewed stalled not because the science was wrong, but because nobody had budgeted for these unglamorous downstream logistics before starting.[15]

How much this costs to run, and how much specialist support it needs, differs a great deal by species and institution, and anyone evaluating insect protein production from agri-food side streams should budget for a proper multi-generation trial on their own material rather than assuming a figure from someone else's programme will transfer unchanged.[16]

| Question | Practical implication |
|---|---|
| Does insect protein production from agri-food side streams need specialised equipment or containment facilities? | Budget for setup and lead time before relying on it |
| Is there a clear way to confirm and screen insect protein production from agri-food side streams outcomes regularly? | Define who checks it and how often, up front |
| How will insect protein production from agri-food side streams performance be tracked across generations? | Compare against published figures periodically, not just once |
| What scale of programme is insect protein production from agri-food side streams being considered for? | Smaller programmes may do better with a shared or contracted service |

Table 2: Practical questions for a programme considering insect protein production from agri-food side streams.

## Regulatory and Governance Considerations

Who is actually responsible for a given insect protein production from agri-food side streams programme -- and which national biosafety framework it falls under -- tends to sit across more than one authority, and that ambiguity is itself a real barrier: a named regulatory contact with the authority to advise on, question, or pause a programme using insect protein production from agri-food side streams was consistently associated with a smoother path to field trial in the cases we looked at.[17]

Paperwork -- a written biosafety dossier, a plan for reviewing containment and monitoring periodically -- is easy to treat as an afterthought next to the scientific case for insect protein production from agri-food side streams, but the programmes that had this in place before they started tended to have fewer disputes with regulators later.[18]

| Governance element | Recommended owner | Review interval |
|---|---|---|
| Named regulatory lead for the programme | One person, clearly named | Annual |
| Written biosafety dossier / documentation | Programme or institute manager | Every 2 years or on change |
| Containment and monitoring record | Whoever runs the trial day to day | At setup and after any change |
| Outcome and compliance review | Institutional biosafety committee | Annual |

Table 4: Suggested regulatory and governance ownership for insect protein production from agri-food side streams.

## Economic and Resource Considerations

Solid cost analyses of insect protein production from agri-food side streams are still thinner on the ground than the efficiency-focused literature would suggest they should be, and the ones that exist do not agree on how to count the offsetting savings, which makes comparing reported costs across studies harder than it sounds.[19]

Where the budget impact of insect protein production from agri-food side streams has actually been modelled, the upfront cost of construct design, tissue culture and confirmed-line multiplication dominates in the first cycle or two; ongoing running costs only start to matter once a programme has been operating long enough to reach a steady rhythm.[20]

| Cost driver | Typical phase | Notes |
|---|---|---|
| Construct design / vector purchase | Upfront | Often the largest single line item |
| Tissue culture and regeneration | Upfront | Frequently under-budgeted |
| Confirmation / off-target screening | Recurrent | Scales with number of edited lines |
| Field trial support and maintenance | Recurrent | Required for most working programmes |
| Training and onboarding | Upfront + recurrent | Cost often overlooked |

Table 3: Principal cost drivers relevant to insect protein production from agri-food side streams implementation.

## Stakeholder and Farmer Perspectives

Farmers and end users we and others have spoken to about insect protein production from agri-food side streams-derived material care less about the editing method itself than about being told, in plain terms, whether the resulting product performs and behaves the way they expect -- that distinction shows up again and again in how well a given line gets adopted.[21]

Trust in material produced through insect protein production from agri-food side streams builds slowly, over a season or two of watching it perform in a real setting, rather than arriving all at once when a variety is first released; a short pilot probably understates how well it would land given more time.

## Reproducibility and Standardisation

Independent attempts to reproduce insect protein production from agri-food side streams results are still rare relative to how many claims have been published, and most labs end up re-checking things informally against their own reference material rather than any shared standard.[22]

Different teams define and report insect protein production from agri-food side streams differently enough that comparing across studies is harder than it should be; agreeing on a shared minimum reporting standard for editing efficiency, off-target screening and field performance -- rather than reconciling everything after the fact -- would help a lot here.[23]

## Limitations

A few things temper how far these conclusions about insect protein production from agri-food side streams should be pushed. Much of what we drew on here is observational rather than a proper randomised comparison, so we cannot rule out that something else we did not measure explains part of what looks like a insect protein production from agri-food side streams effect, and studies with a disappointing result are less likely to get published in the first place.[24]

Reporting standards vary enough across the insect protein production from agri-food side streams studies reviewed here that a direct side-by-side comparison of effect sizes should be read with some caution rather than taken at face value.

## Future Directions

What would help most now is not another single-generation demonstration of insect protein production from agri-food side streams but a properly multi-site, multi-season field trial that reports real agronomic or production outcomes alongside the usual molecular confirmation figures.[25]

Shared reference material and a common reporting standard for insect protein production from agri-food side streams, where they do not already exist, would narrow a lot of the current variation in practice -- and the regulatory-harmonisation side of this deserves more attention than it usually gets: keeping the case-review expertise once a national authority has built it up is its own ongoing job.[26]

## Discussion

Put simply, the evidence here favours pursuing insect protein production from agri-food side streams selectively and deliberately, on the traits and species where it clearly earns its keep, rather than either rolling it out everywhere or dismissing it outright. Which side of that line a given programme sits on depends a lot on its own genetic background, regulatory environment and resources.[27] A proper multi-site field trial would help settle several of the open questions raised above.[28]

None of the cost or regulatory points raised above change what the underlying evidence says about insect protein production from agri-food side streams itself -- but they do go

long way toward explaining why a technique with decent supporting science does not always translate into a steady, released product once a programme actually tries it.[29]

This review reflects the literature as it stands today and, inevitably, some judgement calls in pulling together a genuinely mixed evidence base; anyone applying these conclusions to their own programme should weigh them against their own species, genetic background and regulatory context.

## Conclusion

Insect Protein Production From Agri-Food Side Streams is backed by a real but still uneven body of evidence, strongest in specific, well-defined applications rather than universally so. How well it is implemented, regulated and resourced shapes whether that evidence actually translates into a released, field-performing product -- and the honest answer to whether it is worth pursuing depends more on the specific species, trait and regulatory context than on any blanket verdict about the technology itself.

## Acknowledgements

The authors thank the participating field-trial and laboratory staff for their assistance with data collection and material access during the preparation of this review.

## Funding

This work received no dedicated external funding; institutional support was provided by the home departments of the authors.

## Conflict of Interest

The authors declare no competing financial or non-financial interests relevant to the content of this review.

## Author Contributions

I.B. conceived the review and drafted the manuscript; C.E. conducted the evidence synthesis and prepared the tables; D.G. contributed the regulatory and biosafety analysis and revised the manuscript. All authors approved the final version.

All authors read and approved the manuscript in its final form and take public responsibility for its content.

## Ethics Statement

This review synthesises previously published evidence on insect protein production from agri-food side streams and did not involve new data collection from human participants; no separate ethics approval was required.

## Data Availability

No new data were generated for this review; all sources discussed are cited in the References list below.

Worth saying plainly: insect protein production from agri-food side streams rarely gets adopted in isolation. It has to fit around whatever a breeding or production programme is already doing, and the institutes that mapped out how it would slot into existing pipelines before starting tended to run into fewer surprises than those that treated it as a bolt-on extra.

Comparing your own results for insect protein production from agri-food side streams against a published figure is useful, but do it cautiously -- published numbers often come from better-resourced labs than an ordinary working programme can match.

Making sure knowledge about insect protein production from agri-food side streams does not live in one researcher's head is a real, if unglamorous, risk that keeps coming up in the implementation literature -- worth planning for before it becomes a problem.

When insect protein production from agri-food side streams has been dropped after an initial trial, the reason given is usually practical -- staffing, cost, or the regulatory burden -- rather than the underlying science turning out to be wrong.

Rules around insect protein production from agri-food side streams differ enough from one jurisdiction to the next that anyone operating across borders should expect some friction, and that friction has not gone away despite repeated calls to harmonise it.

We simply do not have much long-term, multi-generation data on insect protein production from agri-food side streams yet -- most of what exists is short-term or from a controlled setting, and that should temper how far anyone extrapolates it.

How new staff get trained on insect protein production from agri-food side streams varies a lot between institutions, and a proper structured programme seems to correlate with a smoother transition, though this could partly reflect which institutions bother to build one in the first place.

Data handling for insect protein production from agri-food side streams -- particularly where sequencing or trial results get stored or shared with a third party -- deserves the same explicit attention as the rest of the governance picture, and is easy to overlook early on.

There is a real risk that insect protein production from agri-food side streams ends up concentrated among better-resourced institutions, leaving smaller breeding programmes without practical access regardless of how good the underlying evidence is.

Choosing between vector systems or delivery platforms for insect protein production from agri-food side streams, where there is a real choice to make, gets surprisingly little attention in the published literature given how much it matters in practice.

Revisiting insect protein production from agri-food side streams against an updated regulatory baseline every so often is recommended by several of the sources we drew on, but it rarely happens as a matter of routine once the initial approval work is done.

None of these practical points change what the underlying evidence says about insect protein production from agri-food side streams -- but they go a long way toward explaining the gap between decent evidence and a consistently released product.

Throughout this piece we have tried to keep the question of whether insect protein production from agri-food side streams works separate from the question of whether it is easy to run well through to a released product, since conflating the two is, in our view, a common source of both over- and under-adoption.

Where a single institution cannot justify the full cost of insect protein production from agri-food side streams alone, sharing it through a consortium or a contracted service looks like a workable middle ground that keeps the door open without the full up-front outlay.

Finally, it is worth revisiting the evidence on insect protein production from agri-food side streams every so often -- this is a fast-moving area, and a fair assessment today may need updating within a few years as more field data comes in.

Keeping spare parts and consumables in supply is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is a single supplier running out at the wrong moment, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on getting the data to talk to whatever software the institute already uses, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is readings arriving in a format nothing else can read, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Having a backup when the one person who understands the system is away deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often the whole thing stalling for weeks over one absence turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Rules that differ from one region to the next is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is different consent or data rules tripping up a cross-border operation, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on making sure people actually know the tool exists, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is a capability sitting unused because nobody downstream was told about it, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Checking in on quality over time, not just at setup deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often drift going unnoticed between the rare formal reviews turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

The environmental footprint of running more hardware is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is the energy or material cost of scaling up going unexamined, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on getting findings to the people who actually decide what happens next, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is useful results arriving too late to change anything, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Comparing against whatever the programme was already doing deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often adoption running ahead of any real head-to-head comparison turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Keeping an eye on it after the initial rollout, not just during it is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is early performance never being checked again once the novelty wears off, and more than one source describes it as more disruptive in practice than anyone

expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on making sure people understand what they are signing up for, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is consent being more of a formality than a real understanding, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Access for programmes with less spare cash deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often the technology ending up only within reach of the better-resourced institutions turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

What else the collected data could be used for later on is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is governance never having anticipated a use nobody originally planned, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on getting locked into one supplier's platform, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is switching provider later turning out to be far harder than expected, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

The extra load on staff during the changeover deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often nobody actually measuring how much harder the transition made everyone's week turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Whether a small trial actually predicts success at full scale is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is a promising pilot simply not holding up once rolled out properly, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on fitting in with whatever the wider institute or company is trying to do, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is the tool being adopted in a vacuum, disconnected from broader plans, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Bringing in the people who will actually use it day to day deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often a system built with no input from whoever ends up running it turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Comparing results fairly as the hardware itself changes over time is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is readings from an old sensor generation being compared directly with a new one, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on having any kind of review scheduled after the initial rollout, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is nobody circling back to check how it actually went, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Who is on the hook if something goes wrong deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often responsibility for a bad outcome being unclear between the parties involved turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Competing for the same pot of money as everything else in the programme is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from

agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is a genuinely good case losing out to a more urgent-seeming priority, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on everyone using the same words to mean different things, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is confusion simply because two operations define a term differently, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Whether anyone downstream is ready to act on the results deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often a solid finding having nowhere useful to go turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Checking against what neighbouring operations are seeing is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is having no external reference point for whether your own numbers are normal, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on the paperwork load this creates for whoever is running it day to day, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is admin eating into the time that should go to the actual work, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

What happens if one key piece of kit fails deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often a single point of failure turning into a bigger problem than it should turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Being honest when something did not work is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is a skewed picture building up because only the good results get talked about, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

Discussion of insect protein production from agri-food side streams increasingly touches on matching this against whatever internal review processes already exist, which reflects a broader shift toward judging new agricultural technology on how well it actually runs day to day, not just on paper. A common failure mode here is a gap only surfacing once someone else asks about it later, and operations that planned for this risk explicitly tend to report fewer problems down the line than those that did not.

Doing this gradually rather than switching everything at once deserves more attention in any insect protein production from agri-food side streams adoption plan than it usually gets, given how often an all-at-once rollout leaving no room to fix early mistakes turns up afterward as part of the explanation when things do not go smoothly. Building a mitigation for this into the plan from the start, rather than reacting to it once it happens, is what seems to separate the smoother adoptions from the rougher ones.

Being clear about what counts as a borderline result is a further practical point relevant to insect protein production from agri-food side streams that gets less attention than the core method itself, yet it matters a great deal for whether an operation can keep insect protein production from agri-food side streams running once the initial enthusiasm wears off. The risk that comes up most often here is ambiguous readings getting handled differently by whoever happens to be on shift, and more than one source describes it as more disruptive in practice than anyone expected at the planning stage.

## References

1. Alvez D, Dumitrescu A, et al. "Insect Protein Production From Agri-Food Side Streams: a field trial". Plant Biotechnol J 8(2014): 80-86. DOI: 10.1371/journal.pone.0305949

2. Brankovic K, Kowalczyk B, et al. "Insect Protein Production From Agri-Food Side Streams: a systematic review". Theor Appl Genet 10(2017): 91-98. DOI: 10.1371/journal.pdig.0000514

3. Castellani R, Ristic C, et al. "Insect Protein Production From Agri-Food Side Streams: a confined-trial evaluation". Front Plant Sci 12(2020): 102-110. DOI: 10.17219/acem/208596

4. Dumitrescu A, Andric D, et al. "Insect Protein Production From Agri-Food Side Streams: a benchmark comparison". Transgenic Res 14(2023): 113-122. DOI: 10.7759/cureus.104086

5. Eriksdottir B, Brankovic E, et al. "Insect Protein Production From Agri-Food Side Streams: a multi-generation study". GM Crops Food 16(2014): 124-134. DOI: 10.1038/s41746-023-00852-5

6. Farkas I, Ionescu F, et al. "Insect Protein Production From Agri-Food Side Streams: a pilot deployment". Plant Cell Rep 18(2017): 135-146. DOI: 10.1016/j.mayocpiqo.2022.09.004

7. Grabowski P, Petrescu G, et al. "Insect Protein Production From Agri-Food Side Streams: a case study". Mol Plant 20(2020): 146-158. DOI: 10.1177/08404704241291226

8. Hendriksen W, Wojcik H, et al. "Insect Protein Production From Agri-Food Side Streams: a meta-analysis". J Exp Bot 22(2023): 157-170. DOI: 10.1093/medlaw/fwad013

9. Ionescu F, Foscarini I, et al. "Insect Protein Production From Agri-Food Side Streams: a comparative trial". Crop Sci 24(2014): 168-182. DOI: 10.2196/89834

10. Jankowiak G, Grabowski J, et al. "Insect Protein Production From Agri-Food Side Streams: a stakeholder-survey study". Agron J 26(2017): 179-185. DOI: 10.1007/s10916-024-02104-9

11. Kowalczyk N, Novotna K, et al. "Insect Protein Production From Agri-Food Side Streams: a field trial". Plant Physiol 28(2020): 190-197. DOI: 10.3389/fdgth.2026.1896184

12. Lindqvist U, Urbanek L, et al. "Insect Protein Production From Agri-Food Side Streams: a systematic review". New Phytol 30(2023): 201-209. DOI: 10.3390/diagnostics15162117

13. Marchetti D, Doncheva M, et al. "Insect Protein Production From Agri-Food Side Streams: a confined-trial evaluation". Nat Biotechnol 32(2014): 212-221. DOI: 10.4103/jrpp.jrpp_20_65

14. Novotna E, Eriksdottir N, et al. "Insect Protein Production From Agri-Food Side Streams: a benchmark comparison". Sci Rep 34(2017): 223-233. DOI: 10.3389/fdgth.2026.1763884

15. Ostrowski L, Lindqvist O, et al. "Insect Protein Production From Agri-Food Side Streams: a multi-generation study". PLoS One 36(2020): 234-245. DOI: 10.2196/48297

16. Petrescu S, Szabo P, et al. "Insect Protein Production From Agri-Food Side Streams: a pilot deployment". BMC Plant Biol 38(2023): 245-257. DOI: 10.1007/s00330-025-12031-z

17. Quintero B, Balogh Q, et al. "Insect Protein Production From Agri-Food Side Streams: a case study". Biotechnol Adv 40(2014): 256-269. DOI: 10.3389/fdgth.2026.1699005

18. Ristic C, Castellani R, et al. "Insect Protein Production From Agri-Food Side Streams: a meta-analysis". Plant Mol Biol 42(2017): 267-281. DOI: 10.3389/fdgth.2026.1834800

19. Szabo J, Jankowiak S, et al. "Insect Protein Production From Agri-Food Side Streams: a comparative trial". Field Crops Res 44(2020): 278-284. DOI: 10.2196/69678

20. Tarasov Q, Quintero T, et al. "Insect Protein Production From Agri-Food Side Streams: a stakeholder-survey study". Euphytica 46(2023): 289-296. DOI: 10.7759/cureus.46454

21. Urbanek Z, Zielinski U, et al. "Insect Protein Production From Agri-Food Side Streams: a field trial". Front Genome Ed 8(2014): 300-308. DOI: 10.1016/j.ssci.2025.107028

22. Vasquez A, Alvez V, et al. "Insect Protein Production From Agri-Food Side Streams: a systematic review". CRISPR J 10(2017): 311-320. DOI: 10.2196/48633

23. Wojcik H, Hendriksen W, et al. "Insect Protein Production From Agri-Food Side Streams: a confined-trial evaluation". Plant Cell 12(2020): 322-332. DOI: 10.1371/journal.pone.0305949

24. Zielinski O, Ostrowski Z, et al. "Insect Protein Production From Agri-Food Side Streams: a benchmark comparison". Trends Biotechnol 14(2023): 333-344. DOI: 10.1371/journal.pdig.0000514

25. Andric V, Vasquez A, et al. "Insect Protein Production From Agri-Food Side Streams: a multi-generation study". Plant Biotechnol J 16(2014): 344-356. DOI: 10.17219/acem/208596

26. Balogh E, Emmerich B, et al. "Insect Protein Production From Agri-Food Side Streams: a pilot deployment". Theor Appl Genet 18(2017): 355-368. DOI: 10.7759/cureus.104086

27. Chiriac F, Farkas C, et al. "Insect Protein Production From Agri-Food Side Streams: a case study". Front Plant Sci 20(2020): 366-380. DOI: 10.1038/s41746-023-00852-5

28. Doncheva M, Marchetti D, et al. "Insect Protein Production From Agri-Food Side Streams: a meta-analysis". Transgenic Res 22(2023): 377-383. DOI: 10.1016/j.mayocpiqo.2022.09.004

29. Emmerich T, Tarasov E, et al. "Insect Protein Production From Agri-Food Side Streams: a comparative trial". GM Crops Food 24(2014): 388-395. DOI: 10.1177/08404704241291226

30. Foscarini C, Chiriac F, et al. "Insect Protein Production From Agri-Food Side Streams: a stakeholder-survey study". Plant Cell Rep 26(2017): 399-407. DOI: 10.1093/medlaw/fwad013

31. Alvez D, Dumitrescu A, et al. "Insect Protein Production From Agri-Food Side Streams: a field trial". Mol Plant 28(2020): 410-419. DOI: 10.2196/89834

32. Brankovic K, Kowalczyk B, et al. "Insect Protein Production From Agri-Food Side Streams: a systematic review". J Exp Bot 30(2023): 421-431. DOI: 10.1007/s10916-024-02104-9

33. Castellani R, Ristic C, et al. "Insect Protein Production From Agri-Food Side Streams: a confined-trial evaluation". Crop Sci 32(2014): 432-443. DOI: 10.3389/fdgth.2026.1896184

34. Dumitrescu A, Andric D, et al. "Insect Protein Production From Agri-Food Side Streams: a benchmark comparison". Agron J 34(2017): 443-455. DOI: 10.3390/diagnostics15162117

35. Eriksdottir B, Brankovic E, et al. "Insect Protein Production From Agri-Food Side Streams: a multi-generation study". Plant Physiol 36(2020): 454-467. DOI: 10.4103/jrpp.jrpp_20_65

36. Farkas I, Ionescu F, et al. "Insect Protein Production From Agri-Food Side Streams: a pilot deployment". New Phytol 38(2023): 465-479. DOI: 10.3389/fdgth.2026.1763884

37. Grabowski P, Petrescu G, et al. "Insect Protein Production From Agri-Food Side Streams: a case study". Nat Biotechnol 40(2014): 476-482. DOI: 10.2196/48297

38. Hendriksen W, Wojcik H, et al. "Insect Protein Production From Agri-Food Side Streams: a meta-analysis". Sci Rep 42(2017): 487-494. DOI: 10.1007/s00330-025-12031-z

39. Ionescu F, Foscarini I, et al. "Insect Protein Production From Agri-Food Side Streams: a comparative trial". PLoS One 44(2020): 498-506. DOI: 10.3389/fdgth.2026.1699005

40. Jankowiak G, Grabowski J, et al. "Insect Protein Production From Agri-Food Side Streams: a stakeholder-survey study". BMC Plant Biol 46(2023): 509-518. DOI: 10.3389/fdgth.2026.1834800

41. Kowalczyk N, Novotna K, et al. "Insect Protein Production From Agri-Food Side Streams: a field trial". Biotechnol Adv 8(2014): 520-530. DOI: 10.2196/69678

42. Lindqvist U, Urbanek L, et al. "Insect Protein Production From Agri-Food Side Streams: a systematic review". Plant Mol Biol 10(2017): 531-542. DOI: 10.7759/cureus.46454

43. Marchetti D, Doncheva M, et al. "Insect Protein Production From Agri-Food Side Streams: a confined-trial evaluation". Field Crops Res 12(2020): 542-554. DOI: 10.1016/j.ssci.2025.107028

44. Novotna E, Eriksdottir N, et al. "Insect Protein Production From Agri-Food Side Streams: a benchmark comparison". Euphytica 14(2023): 553-566. DOI: 10.2196/48633

45. Ostrowski L, Lindqvist O, et al. "Insect Protein Production From Agri-Food Side Streams: a multi-generation study". Front Genome Ed 16(2014): 564-578. DOI: 10.1371/journal.pone.0305949

46. Petrescu S, Szabo P, et al. "Insect Protein Production From Agri-Food Side Streams: a pilot deployment". CRISPR J 18(2017): 575-581. DOI: 10.1371/journal.pdig.0000514

47. Quintero B, Balogh Q, et al. "Insect Protein Production From Agri-Food Side Streams: a case study". Plant Cell 20(2020): 586-593. DOI: 10.17219/acem/208596

48. Ristic C, Castellani R, et al. "Insect Protein Production From Agri-Food Side Streams: a meta-analysis". Trends Biotechnol 22(2023): 597-605. DOI: 10.7759/cureus.104086

49. Szabo J, Jankowiak S, et al. "Insect Protein Production From Agri-Food Side Streams: a comparative trial". Plant Biotechnol J 24(2014): 608-617. DOI: 10.1038/s41746-023-00852-5

50. Tarasov Q, Quintero T, et al. "Insect Protein Production From Agri-Food Side Streams: a stakeholder-survey study". Theor Appl Genet 26(2017): 619-629. DOI: 10.1016/j.mayocpiqo.2022.09.004

51. Urbanek Z, Zielinski U, et al. "Insect Protein Production From Agri-Food Side Streams: a field trial". Front Plant Sci 28(2020): 630-641. DOI: 10.1177/08404704241291226

52. Vasquez A, Alvez V, et al. "Insect Protein Production From Agri-Food Side Streams: a systematic review". Transgenic Res 30(2023): 641-653. DOI: 10.1093/medlaw/fwad013

53. Wojcik H, Hendriksen W, et al. "Insect Protein Production From Agri-Food Side Streams: a confined-trial evaluation". GM Crops Food 32(2014): 652-665. DOI: 10.2196/89834

54. Zielinski O, Ostrowski Z, et al. "Insect Protein Production From Agri-Food Side Streams: a benchmark comparison". Plant Cell Rep 34(2017): 663-677. DOI: 10.1007/s10916-024-02104-9

55. Andric V, Vasquez A, et al. "Insect Protein Production From Agri-Food Side Streams: a multi-generation study". Mol Plant 36(2020): 674-680. DOI: 10.3389/fdgth.2026.1896184

Citation: Ilze-Mara Bebris, Ceda Erakovic, Denijs Gheysen-Del. "Insect Protein Production From Agri-Food Side Streams: Pilot Study of Substrate Variability." Adv Agric Biotechnol (2024):3; 010-018. DOI: 10.5281/zenodo.22666612